Managing sore throat in the pharmacy

Learning objectives:

Refresh knowledge of the causes and symptoms of sore

throat

f d sore Understand how benzydamine hydrochloride works Be able to recommend an appropriate product to manage sore throat.

Difflam

Sore throats are common and are usually caused by a viral infection of the upper respiratory tract, such as a cold or influenza. Very occasionally they can be caused by a bacterial infection. Sore throats can also be caused by laryngitis, tonsillitis, strep throat (a bacterial throat infection) or glandular fever.¹ The highest incidence of sore throat is in children during the winter months.²

A sore throat due to a viral or bacterial cause is usually a self-limiting condition; symptoms resolve within one week in 85% of people.³ Customers should be advised to see their GP if their symptoms have not improved after one week.⁴

Symptoms of sore throat include a painful throat, especially when swallowing; a dry, scratchy throat; and redness in the back of the mouth. Other symptoms include bad breath, a mild cough and swollen neck glands. Symptoms are similar for children, but they can also have a temperature and appear less active.¹

Job number: DIFF-2024-0003 Date of preparation: February 2024.



Managing **sore throat**

NICE recommends that customers presenting with sore throat should be advised that:5



Adequate fluid should be taken during the course of the illness to keep the throat moist and prevent dehydration



Medicated lozenges

containing a local anaesthetic, non-steroidal anti-inflammatory drug (NSAID) or an antiseptic agent may provide temporary relief from throat pain.



Ibuprofen or paracetamol can be used to help relieve the pain and discomfort of a sore throat as well as reducing fever.

Benzydamine hydrochloride

Benzydamine hydrochloride is an NSAID that has a different mechanism of action from other aspirin-like NSAIDs, providing triple action relief from sore throat.



When to **refer**

Most sore throats can be managed with advice and product recommendations from the pharmacy team. However, it is important to know when customers should be referred to their GP.

WWHAM	When to refer
W	 Customer has a weakened immune system or is undergoing chemotherapy¹ Note that benzydamine can very rarely cause bronchospasm so should be used with caution by customers with asthma⁶
	Customer appears significantly unwell
W	 Customer gets sore throats frequently
	 Signs of a more serious medical condition
	 Has high risk of complications^{1,5}
H	 Customer has had symptoms for longer than one week with no improvement⁴
A	 Customer has tried pain relief or medicated lozenges/ sprays, and seen no improvement after three days⁵
M	 On any other medication; look out for carbimazole, methotrexate or other DMARDs (disease-modifying anti- rheumatic drugs)⁵

References: 1. https://www.nhs.uk/conditions/sore-throat/ 2. https://cks.nice.org.uk/topics/sore-throat-acute/backgroundinformation/prevalence/ 3. https://cks.nice.org.uk/topics/sore-throat-acute/background-information/prognosis/ 4. Duerden, M., Benkinsopp, A. and Blenkinsopp, J. (2018). Symptoms in the Pharmacy (8th Edition) 5. https://cks.nice.org.uk/topics/sore-throatacute/management/management/ 6. https://www.medicines.org.uk/emc/product/827/smpc/print

Online references last accessed February 2024.

Job number: DIFF-2024-0003. Date of preparation: February 2024.

The **Difflam** product range

Difflam has a range of products that you can recommend to customers based on their personal preference.

Difflam Sprav

- Fast acting, long lasting relief of sore throat pain
- Triple action: Anti-inflammatory. Analgesic. Anaesthetic
- Directly targets the site of pain
- Suitable for adults and children of all ages
- Sugar, lactose and gluten free

Difflam Sore Throat Rinse

- Fast acting, long lasting relief of sore throat pain
- Triple action: Anti-inflammatory, Analgesic, Anaesthetic
- Includes measuring cup for accurate dosing
- Suitable for adults and children over 12 years old
- Sugar, lactose and gluten free



Difflam 3mg Lozenges are also available

Benzydamine hydrochloride 3mg | P

ESSENTIAL INFORMATION:

ESSENTIAL INFORMATION: Difflam™ 0.15 % w/v Spray (benzydamine hydrochloride). Presentation: Difflam™ Spray contains benzydamine hydrochloride 0.15% w/v. Each metered dose pump spray delivers approximately 175mcl per puff. Indications: Difflam™ Spray is a locally acting analgesic and anti-inflammatory treatment for the throat and mouth. It is especially useful for the relief of pain in traumatic conditions such as following surgical removal of tonsils or the use of a naso-gastric tube; dental surgery. Dosage and administration: Adults and elderly. 4 to 8 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children and eff. 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-10: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-10: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. Children acded 5-12: 4 puffs, 1½ to 3 hourly. PL 46302/0160 Legal category: P.

Difflam™ Sore Throat Rinse & Difflam™ Oral Rinse (benzydamine hydrochloride). Presentation: Difflam[™] Sore Throat Rinse & Difflam™ Oral Rinse contain benzydamine hydrochloride 0.15% w/v. Presented as a gargle/oral rinse in a 200ml and 300ml glass bottle respectively. Indications: Locally acting analgesic and anti-inflammatory treatment for the relief of painful inflammatory conditions of the throat including pharyngitis. Dosage and administration: Rinse or gargle and anti-inflammatory treatment for the relief of painful inflammatory conditions of the throat including pharyngitis. Dosage and administration: Rinse or gargle with T5ml (using measuring cup provided) every 1½-3 hours as required. Expel solution from the mouth after use. If stinging occurs the rinse may be diluted with water. Uninterrupted treatment should not exceed seven days, except under medical supervision. It is not suitable for children aged 12 years or under. **Contra-indications**: Patients with hypersensitivity to ingredients. **Warnings and Precautions**: Not advisable in patients with hypersensitivity to acetylsalicylic acid or other NSAIDS. Bronchospasm may be precipitated in patients suffering from or with a previous history of bronchial asthma. Caution should be excertised in these patients. The small amount of alcohol in this medicine will not have any noticeable effects. Some inactive ingredients may cause allergic reactions. Avoid contact with the eyes. Interactions: None known. **Pregnancy and lactation**: DifflamTM Sore Throat Rinse should not be used in pregnancy or lactation unless considered essential by the physician. There is no evidence of a teratogenic effect in animal studies. **Undesirable effects: Serious (frequency not known)**: Anaphylactic reaction which can be potentially life-threatening. Hypersensitivity reactions, Angioedema (**Uncommon**). Oral tissue numbness or 'stinging' sensations in the mouth may occur. (**Very rare)**: Privritia, photosensitivity reaction, layngespasm/hornchospasm. Prescribers should consult the Summary of Product **Licence number**: DifflamTM Sore Throat Rinse 200ml bottle £8.99, DifflamTM Oral Rinse 800ml bottle £11.19 **Product Licence number**: DifflamTM Sore Throat Rinse PL 46302/0154 & DifflamTM Oral Rinse PL 46302/0159 **Legal category**: P

Marketing Authorisation Holder: Mylan Products Ltd, Station Close, Potters Bar, Hertfordshire, EN6 1TL, United Kingdom. Date of preparation of Essential information: August 2023 Ref number: DIFF-2023-0038

Please continue to report suspected adverse drug reactions with any medicine or vaccine to the MHRA through the Yellow Card Scheme. It is easiest and quickest to report adverse drug reactions online via the Yellow Card Scheme website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. Alternatively, you can report via some clinical IT systems (EMIS/SystemOne/Vision/MiDiatabank) or by calling the Commission on Human Medicines (CHM) free phone line: 0800-731-6789. Adverse reactions/events should also be reported to MAH at e-mail address: pv.uk@viatris.com



Benzydamine

hydrochloride 0.15% w/v P



Sore Throat Rinse

Benzydamine

Difflam and complete the e-module



Anti-inflammaton